## PROBLEM OF THE MONTH OCTOBER 2014 - SOLUTION

Determine:

$$\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{3n-2} = 1 - \frac{1}{4} + \frac{1}{7} - \frac{1}{10} + \cdots$$

Solution:

$$\frac{\ln(2)}{3} + \frac{\pi}{3^{\frac{3}{2}}}.$$

You may recall the formula for a finite geometric series which says for  $x \neq 1$ :

$$1 + x + x^{2} + x^{3} + \dots + x^{n-1} = \frac{1 - x^{n}}{1 - x}.$$

Replacing x by  $-x^3$  gives for  $x \neq 1$ :

$$1 - x^{3} + x^{6} - x^{9} + \dots + (-1)^{n+1} x^{3n-3} = \frac{1 - (-1)^{n} x^{3n}}{1 + x^{3}}.$$

Integrating on [0, 1] gives:

$$1 - \frac{1}{4} + \frac{1}{7} - \frac{1}{10} + \dots + \frac{(-1)^{n+1}}{3n-2} = \int_0^1 \frac{1 - (-1)^n x^{3n}}{1 + x^3} \, dx = \int_0^1 \frac{1}{1 + x^3} \, dx + \int_0^1 \frac{(-1)^n x^{3n}}{1 + x^3} \, dx.$$
(1)

Next we observe since  $0 \le x \le 1$  that:

$$\left| \int_0^1 \frac{(-1)^n x^{3n}}{1+x^3} \, dx \right| \le \int_0^1 \frac{x^{3n}}{1+x^3} \, dx \le \int_0^1 x^{3n} \, dx = \frac{1}{3n+1} \to 0 \text{ as } n \to \infty.$$

Thus we see by taking limits in (1) that:

$$\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{3n-2} = \int_0^1 \frac{1}{1+x^3} \, dx.$$

Now using partial fractions:

$$\int_0^1 \frac{1}{1+x^3} dx = \frac{1}{3} \int_0^1 \left( \frac{1}{1+x} + \frac{-x+2}{x^2-x+1} \right) dx$$
$$= \frac{1}{3} \left( \ln(1+x) - \frac{1}{2} \ln(x^2-x+1) + \sqrt{3} \tan^{-1}(\frac{2}{\sqrt{3}}(x-\frac{1}{2})) \right) |_0^1$$
$$= \frac{\ln(2)}{3} + \frac{2\sqrt{3}}{3} \tan^{-1}(\frac{1}{\sqrt{3}}) = \frac{\ln(2)}{3} + \frac{\pi}{3^{\frac{3}{2}}}.$$